

Effect of Saturated Solution of Potassium Chloride, Copper Sulphate, Ammonium Molybdate on a Plant Growth Promoting Rhizobacteria - *Azotobacter Chroococcum*.

Deokar. S.C.

Department of Chemistry, K.V.N Naik College, Nasik - 422002.

Abstract:

The nitrogen assimilating bacterium *Azotobacter chroococcum* is a non symbiotic nodule forming bacterium which grows in the Jenson media is devoid of nitrogen ingredients, here saturated solution of Potassium chloride (KCl), Copper sulphate (CuSO_4), Ammonium molybdate (NH_4MoO_4) were separately used. The bacterium *Azotobacter chroococcum* was inoculated in the Jenson broth and incubated for 48 hrs. 1ml of this incubated broth was serially diluted to 10^9 dilution and 0.1 ml of 10^9 dilution was spread inoculated on Jenson agar media plate with thickness of 5mm, single sterile whatman filter paper disc grade 1 was dipped in sterile saturated solution of compound and placed in the centre of spread inoculate media in petri plate. The petri plate was kept in fridge at 4°C for 5 minutes for diffusion. The petri plate was incubated at 32°C , room temperature for 72 hours. In vitro experimental observations for of KCl, showed bacterial inhibition upto 3 mm radial zone from periphery of the filter paper disc whereas in case of CuSO_4 , NH_4MoO_4 no bacterial inhibition was seen surrounding the periphery of the filter paper disc. These compounds are used in the farming as inorganic input (micronutrient) for agricultural crops.

Keywords : Jenson Broth Media, Jenson Agar Media, PGPR (Plant Growth Promoting Rhizobacteria). Serial Dilution, Zone Of Inhibition, Agriculturally Important Bacteria, Saturated Solution.

Introduction

Agricultural crops yields are dependent on soil fertility, thus the difference between survival and extinction for most land-based life is characterized by the thin layer of soil covering earth's surface (Doran and Zeiss, 2000). Soil is the living mixture of minerals and organisms that provides vital nutrient and a healthy environment that nurtures crops growth, therefore the soil is divided into two parts as biotic and abiotic (living and non living).

Fertiliser is any substance to add nutrient to the soil to promote soil fertility and increase plant growth. Soil health relies on the balance of macronutrient as well as microbial health. The

scientists concludes that 50mM of ionic strength is required for adherence of microbial mass to the sand surface Saeed Torkzaban et al, Shiva S. Tazehkhand et al, Sharon L. Walker et al and Scott A. Bradford et al, 5 April 2008 (Transport and fate of bacterial in porous media coupled effects of chemical conditions and pore space geometry, Water resources research, Vol 44, W04403, doi: 10.1029/2007, WR006541, 2008).

Organic and inorganic fertilizers continuously applied for 59 years on farm soil (clay loam, orthic luvisol) studies where the soil applications treatments consisting the combinations as manure + NPK, compost + NPK, cattle manure + straw + NPK were compared with cattle slurry + straw + NPK revealed that, the cattle slurry + straw was most favorable in increasing total C, N, hot water soluble C, microbial biomass C, and dehydrogenase activity. (Influence of long term application of organic and inorganic fertilizer on soil properties, T. Simon et al, A.Czako et al, Plant soil environment.Vol. 60. 2014, No. 7: 314-319. Plant use 50% of nitrogenous fertilizer whereas 2 – 20% lost on evaporation, 15 -25% react with organic compounds and 2-10% interfere surface and ground water, apart decrease in pH of soil to critical level along with air pollution by nitrogen oxide (NO, N₂O, NO₂) which increases from 0.2-0.3% each year, leading in nitrate content threatening human life (Serpil Savei et al “An agricultural pollutant: Chemical fertilizer”, International journal of environmental science and development, Vol. 3, No. 1, Feb 2012).

The optimum concentration of a-biotic entities and N₂-fixing microbes are directly influenced by “rhizospheric” agriculturally important microorganisms (Jonas *et al.*, 2011). Therefore ideal microcosms is needed for agriculture helping microorganisms to dwell, and this favor the commercial crop yields through fulfilling their own nutrient requirements. The practice of broadcasting of inorganic fertilizers through manually or machines by farmers is always more than scientifically recommended. During irrigation such granules in soil gets solubilised which directly affect the microbial population in its vicinity. The study of fate of *Azotobacter chroococcum* a potential PGPR (Plant Growth Promoting Rhizobacteria) is therefore experimented in-vitro for Ferrous and Boron through Ferrous sulphate and Boric acid.

The production of bio-inoculants, PGPR (Plant Growth Promoting Rhizobacteria) are grown in aseptic conditions in desired media which are then mixed with suitable inert carriers

such as lignite. These bio-inoculants so produced provide nutritional support to commercial crops (Gomare, *et al.*, 2013). The present study is made to find the effect of saturated compound of plant micronutrient on *Azotobacter chroococcum* bio-inoculant which is wide-spread used for all monocot crop.

Materials and Methods

Sample Collection

Bacterial Strain: *Azotobacter chroococcum* are the native non nodule forming bacteria found in nature. The bacterium used in the present study is *Azotobacter chroococcum* which is isolated from Biofertilizer Packet produced by M.P Agro Ind., Bhopal, these strains are said to have high nitrogen fixing efficiency.

a) Preparation of saturated solution –

The saturated solution of of KCl, CuSO₄, NH₄MoO₄ is prepared by gradually adding small quantities of compound in 100 ml of double distill water in 500 ml beaker. The stirring is facilitated by using magnetic stirrer till no more compound gets solubilised. The solution is poured in 250 ml conical flask and stoppered with cotton plug and steam sterilized.

b) Preparation of broth culture of Bacterial strain :-

The pure culture of *Azotobacter chroococcum* are inoculated in Jenson broth media and incubated at room temperature for 48 hours to 72 hours till the cell concentration exceeds the optical density (OD) 1 at 620 nm and a viable cell count of 1.0×10^9 per ml of matured (stationary phase bacteria) broth. This matured broth is then diluted to 10^9 and its 0.1 ml is used as inoculum on experimental Jenson agar plate medium, inoculums is spread evenly with the help of sterile spreader.

c) Preparation of filter paper discs :-

Whatman filter paper grade 1 is evenly punched with help of punching machine and several uniform discs were prepared. These discs were wrapped in brown paper and then steam sterilized.

d) Studies of Saturated Concentration of KCl, CuSO₄, NH₄MoO₄ on *Azotobacter chroococcum*:-

Aseptically pre-sterilised whatman filter paper grade 1 disc were dipped in saturated solution and placed in the centre of freshly inoculated *Azotobacter chroococcum* and spread plated on petri plate of the Jenson agar medium. Separate experiments were performed for Ferrous sulphate and Boric acid. The plates were kept in 4°C in refrigerator for diffusion and then uprightly incubated at 32°C, room temperature for 72 hours.

Results and Discussion.

Effect of Saturated solution of KCl, CuSO₄, NH₄MoO₄ on *Azotobacter chroococcum* :-

Zone of inhibition of growth of *Azotobacter chroococcum* is observed in the experiment conducted for KCl. The experiment with KCl showed no growth of bacterium at 3mm as zone radius, whereas growth was seen surrounding the saturated filter disc in experiment using CuSO₄ and NH₄MoO₄. (Table 1). Trace elements/metal function as co-factors in enzymatic reactions, stabilizing structure of enzyme itself. (Zhuoer Lin *et al.*, 2009).

The chief objective of the present investigation is to know the individual effect of concentration of compound of such plant micronutrients on *Azotobacter chroococcum* enabling the fate of useful agriculture microbe on widespread use of granular inorganic fertilizer on farm soil.

Conclusion.

The compounds of micronutrients are well known to impart healthy effect on the plant growth as they are the prime requirement (trace elements) for various vital activities of plants. As such the usual practice adopted by most farmers during broadcasting of inorganic micronutrient is always more than per hectare dose as recommended by agriculture scientists. The in-vitro experimental results concludes the ill effect of such inorganic fertiliser on the population of agriculturally important plant growth promoting rhizobacteria that is *Azotobacter chroococcum*, therefore in this study suggest that inorganic **KCl** of granular or in liquid form may be used in other appropriate

forms whereas CuSO_4 and NH_4MoO_4 do not disturb the existence of Azotobacter chroococcum and can be used through broadcasting for the farming process.

References

- Ahmed I. and Fujiwara T (2010).** Mechanism of Boron Tolerance in Soil Bacteria. *Canadian J Microbiol* **56(1)**: 22–26
- Blevins DG, and Lukaszewski KM (1994).** Proposed Physiologic Functions of Boron in Plants Pertinent to Animal and Human Metabolism. *Environ Health Perspect* **102(Suppl 7)**: 31–33.
- Camacho-Cristóbal JJ, Rexach J, and Fontes AG (2008).** Boron in Plants: Deficiency and Toxicity. *Journal of Integrative Plant Biolo* **50(10)**:1247–1255
- Doran, J.W., and Zeiss, M.R., (2000).** Soil health and sustainability: managing the biotic component of soil quality. *Applied Soil Ecology*. **15**: 3–11
- Gomare, K.S., Mese, M., and Shetkar, Y. (2013)** Isolation of *Rhizobium* and Cost Effective Production of Biofertilizer. *Indian Journal of Life Science*. **2(2)**: 49-53
- H.Babich and G.Stotzky (1978).** Toxicity of zinc to fungi, bacteria and coliphages colon influence of chloride ions. *Applied and environmental microbiology*. **36(6)**: 906-914.
- Jonas. Chianu, E. Nkonya, F. Mairura, Justina. Chianu, F. Akinnifesi (2011).** Biological Nitrogen Fixation and Socioeconomic Factors for Legume Production in Sub-Saharan Africa: a review. *Agronomy for Sustainable Development, Springer Verlag (Germany)*. **31 (1)**: 139-154.
- Khin S. Aye et al (2011),** Investigation on the effectiveness of zinc sulphate and biofertiliser on mustard plant, *World Academy of Science, Engineering and Technology*. 5-2011-03-26, p-233-235
- Singh, B.D. (2008).** Biotechnology, Expanding Horizon. *Kalyani Publishers, New Delhi*. pp 598-603.
- Saeed Torkzaban, Shiva S. Tazehkand, Sharon L. Walker, and Scott A. Bradford. (2008).** Transport and fate of bacteria in porous media: coupled effect of chemical conditions and pore space geometry. *Water resources research*. **44-2008** -wo4403
- Tilak, K.V.B.R (1998).** Bacterial Fertilizer. *Indian council of Agricultural Research, P.I Division, Krishi Anusandhan Bhavan, New Delhi*. pp 4-33.

Tubeck, S., Grzanka, P., Tubeck, I. (2008) Role of Zinc in Hemostasis: A Review. *Biol. Trace Elem. Res.* **121 (1):** 1-8

Zhuoer Lin, Robledo, J.A.F., Cellier, M.F.M., and Vasta, G.R. (2009).Metals and Membrane Metal Transporters in Biological Systems: The Role(s) of Nramp in Host-Parasite Interactions. *The Journal of the Argentine Chemical Society.* **97(1):** 210-225.

Tables

Table 1:- Effect of saturated solution of compounds on *Azotobacter chroococcum*.

S.no	Saturated solution (compound)	Zone of inhibition of <i>Azotobacter chroococcum</i>	
		Radius of no growth	Diameter of no growth
1	KCl	3mm	6mm
2	CuSO ₄	Nil	Nil
3	NH ₄ MoO ₄	Nil	Nil

Note :- The thickness of Jenson agar media in petri plate is 5 mm.