

Reference:

- Abdel-Rahman, A.M., and Martin, 1990.**, The Mount Gharib A-type granite, Nubian Shield: petrogenesis and role of metasomatism at the source. *Contributions to Mineral. Petrol.* 104, 173-183.
- Amin, M. S., Moustafa, G.A. and Zaatout, M.A., 1954**, Geology of Abu Diab District. *Geol. Surv. Egypt*.
- Bentor, Y.K., 1985**: The crustal evolution of the Arabo-Nubian Massif, with special reference to the Sinai Peninsula. *Pre-Cambrian. Res.* V. 28, 1-74p.
- Blott, S.J. and Pye, K. (2001)** GRADISTAT: a grain size distribution and statistics package for the analysis of unconsolidated sediments. *Earth Surface Processes and Landforms* 26, 1237-1248.
- Eby, N.G., 1992**. Chemical subdivision of the A-type granitoids: petrogenetic and tectonic implications. *Geol* 20, 641-644.
- El Kady, A.M., 2014**. (in press) Applied geochemical and mineralogical studied on the granites of Homrit Waggat Central Eastern Desert, Egypt. *Al-Azhar. Univ.* pp155.
- Folk, R.L., 1954**. The distinction between grain size and mineral composition in sedimentary rock nomenclature. *Jour. Geol.* vol. 62, pp. 1–38.
- Folk, R.L., 1980**. Petrology of sedimentary rocks. *hemphill pub. Co., Austin, Texas*, 182p.
- Folk, R.L., Ward, W.C., 1957**. Brazos River bar: a study in the significance of grain size parameters. *Jour. Sed. Petrol* 27, 3–26.
- Friedman, G.M., 1967**. Dynamic processes and statistical parameters compared for size frequency distribution of beach and river. *Jour. Sed. Petrol.*, V. 37. PP. 327-354.
- Hassanen, M.A., 1997**. Post-collision, A-type granites of Homrit Waggat complex, Egypt: petrological and geochemical constraints on its origin. *Precambrian. Res.*, 82, 211-236
- Lumbers, S.B., Wu, T.-W., Heaman, L.M., Vertolli, V.M., and MacRae, N.D., 1991**. Petrology and age of the A-type Mulock granite batholith, northern Grenville Province, Ontario. *Precambrian. Res* 53, 199-231.
- Moghazi, A.M., Mohamed, F.H., El-Sayed, M.M., Kanisawa, S., 1999**. Geochemistry and petrogenesis of late Proterozoic plutonic rock suites in the Homrit Waggat and El-Yatima areas, Eastern Egypt. In *Fourth International Conference on Geochemistry*, Alexandria Univ. Egypt, pp. 1–21.
- Mohamed, F.H., 1993**. Rare metal-bearing and barren granites, Eastern Desert of Egypt: geochemical characterization and metallogenic aspects. *Jour. Afr. Earth. Sci.*, 17, 525-539.
- Moiola, R.J., Weiser, D., 1968**. Textural parameters, an evolution, *Jour. Sed. Petrol.* V. 48, PP. 45-53.
- Niazy, A.E., A. El-Bakri and M.I. Gharib, 1992**. Size analysis and depositional environment of the Nile Valley Quaternary sediments, Northern Upper Egypt, *Sci. Bull. Mania Univ.* 5 (2): 1-29.
- Stern, R.J., Gottfried, D., 1986**. Petrogenesis of late Precambrian (575-600 Ma) bimodal suite in northeast Africa. *Contributions to Mineral. Petrol* 92, 492-501.
- Sylvester, P.J., 1989**. Post-collisional alkaline granites. *J. Geol.* 97, 261-280.